


Date:

/ /



لمجان الدفعات



تفريغ إحصاء صيدلاني

lecture 13

موضوع المحاضرة:

آخى صوصوع بالسكند

رقم المحاضرة:

نور مصطفى

إعداد الصيدلانيه:




Approximating a Binomial Distribution with a Normal Distribution

lecture 13

Normal Approximation of Binomial Distribution

1. Useful because not all binomial tables exist
2. Requires large sample size
3. Gives approximate probability only
4. Need correction for continuity

$$n = 10 \quad p = 0.50$$



Normal Approximation to Binomial Distributions

- The normal distribution is used to approximate the binomial distribution when it would be impractical to use the binomial distribution to find probability.
- Theorem: Normal approximation to a binomial distribution

If $np > 5$ and $n(1-p) > 5$, then the binomial random variable X is approximately normally distributed with mean (μ) and standard deviation (σ) given as follows:

$$\mu = np,$$

$$\sigma = \sqrt{npq}$$

Where n is the number of trial, p is the probability of success on each trial and q is the probability of failure on each trial = $1-p$

Normal Approximation to Binomial Distributions

discrete data ←

Example

For each of the following cases, decide whether you can use the normal distribution to approximate X. if you can find the mean and standard deviation:

$$n = 65, p = 0.51, q = 0.49$$

$$np = (65)(0.51) = 33.15 \geq 5$$

$$nq = (65)(0.49) = 31.85 \geq 5$$

صحيح نشوف اذا بقدرنا تست approximate اولاً
بحسب np و nq

So we can use the normal approximation

$$\text{Mean: } \mu = np = 33.15, \quad \sigma = \sqrt{npq} = 4.03$$

- $N = 15, p = 0.15, q = 0.85$

$$np = (15)(0.15) = 2.25 < 5, \quad nq = (15)(0.85) = 12.75 > 5$$

We cannot use the normal approximation because $np < 5$.

Continuity Correction

← يتعلق $\frac{1}{2}$ من الصغر صغراً
ويزيد $\frac{1}{2}$ من الأكبر كبراً ...

→ على جعل
approximation

The binomial distribution is discrete and to calculate exact binomial probabilities, the binomial formula is used for each value of x . in order for a continuous distribution (like the normal) to be used to approximate a discrete one (like the binomial), a continuity correction should be used. There are two major reasons to employ such a corrections as follows:

* قراءة *


- First, recall that a discrete random variable (like the binomial) can only take on only specified values, whereas a continuous random variable used to approximate it can take on any values whatsoever.

within an interval around those specified values. Hence, when using the normal distribution to approximate the binomial distribution, more accurate approximations are likely to be obtained if a continuity correction is used.

Second, recall that with a continuous distribution (such as the normal), the probability of obtaining a particular value of a random variable is zero. On the other hand, when the normal approximation is used to approximate a discrete distribution, a continuity correction can be employed so that we can approximate the probability of a specific value of the discrete distribution (such as the binomial).

Therefore, when we use a normal distribution which is continuous to approximate a binomial probability which is discrete, we need to move 0.5 unit to the left or to the right of the x -value to include all possible x -values in the interval by using what is called continuity correction. To use a continuity correction to convert the binomial distribution x -values to a normal distribution x -values, we use the following rules:.

Why Probability Is Approximate



Binomial Probability:
Bar Height


Normal Probability: Area Under Curve from 3.5 to 4.5

Continuity Correction $\frac{1}{2}$ \rightarrow 4 \leftarrow $\frac{1}{2}$ \rightarrow 4.5

Correction for Continuity

Continuity Correction

1. A 1/2 unit adjustment to discrete variable
2. Used when approximating a discrete distribution with a continuous distribution
3. Improves accuracy



Using a Normal Distribution to Approximate Binomial Probabilities

1. Determine n and p for the binomial distribution, then calculate the interval:

$$\mu \pm 3\sigma = np \pm 3\sqrt{np(1-p)}$$

(Note: In the original image, 'np' is labeled 'صَدْرَة' and '1-p' is labeled 'q'.)

فاندرها هي معرفة القيمة اذا
كانت تقع من 0 الى N
اذا كان الجواب نعم بنحكي:

If interval lies in the range 0 to n , the normal distribution will provide a reasonable approximation to the probabilities of most binomial events.

Using a Normal Distribution to Approximate Binomial Probabilities

2. Express the binomial probability to be approximated by the form

$$P(x \leq a) \text{ or } P(x \leq b) - P(x \leq a)$$

For example,

الأعداد الأصغر من 3 و 2 و 3

$$P(x < 3) = P(x \leq 2)$$

$$P(x \geq 5) = 1 - P(x \leq 4)$$


$$P(7 \leq x \leq 10) = P(x \leq 10) - P(x \leq 6)$$

Using a Normal Distribution to Approximate Binomial Probabilities

3. For each value of interest a , the correction for continuity is $(a + .5)$, and the corresponding standard normal z -value is


links w/ \hat{p} \leftarrow

$$z = \frac{(a + .5) - \mu}{\sigma}$$



Using a Normal Distribution to Approximate Binomial Probabilities


4. Sketch the approximating normal distribution and shade the area corresponding to the event of interest. Using z score table and the z-value (step 3), To find the shaded area. This is the approximate probability of the binomial event.



Example

Normal Approximation Example

What is the normal approximation of $p(x = 4)$ given $n = 10$, and $p = 0.5$? * حل المسألة تقريبا $n \leftarrow 10$ $p \leftarrow 0.5$



Normal Approximation Solution

1. Calculate the interval:

$$\begin{aligned} np \pm 3\sqrt{np(1-p)} &= 10(0.5) \pm 3\sqrt{10(0.5)(1-0.5)} \\ &= 5 \pm 4.74 = (0.26, 9.74) \end{aligned}$$

Interval lies in range 0 to 10, so normal approximation can be used

✓ تقع في 0 ← 10

2. Express binomial probability in form:

$$P(x = 4) = P(x \leq 4) - P(x \leq 3)$$

هو عينا
إذا كانت x فالجواب صفر
مباشرة...

Normal Approximation Solution


3. Compute standard normal z values: z_{value} & x_{value} \rightarrow μ

$$z = \frac{x - \mu}{\sigma}$$
$$z \approx \frac{(a - .5) - \overbrace{n \cdot p}^{\text{mean } \mu}}{\sqrt{n \cdot p(1-p)}} = \frac{3.5 - 10(.5)}{\sqrt{10(.5)(1-.5)}} = -0.95$$

$$z \approx \frac{(a + .5) - n \cdot p}{\sqrt{n \cdot p(1-p)}} = \frac{4.5 - 10(.5)}{\sqrt{10(.5)(1-.5)}} = -0.32$$

Normal Approximation Solution

4. Sketch the approximate normal distribution:



Normal Approximation Solution

5. The exact probability from the binomial formula is 0.2051 (versus .2034)

