



PHYSIOLOGY

FACULTY OF PHARMACEUTICAL SCIENCES

DR. AMJAAD ZUHIER ALROSAN

LECTURE 15: GASTROINTESTINAL SYSTEM

Objectives

الصورة العامة

<https://youtu.be/eSGRpPcJGIA?si=QWlv1wTiGUgDQxnd>



قبل التفاصيل

1. Discuss **the digestive system.**
2. Describe **neural innervation of the GI tract.**
3. Explore **gastrointestinal motility,** **chewing** and **swallowing.**

(Pages 886- 901 of the reference)

THE DIGESTIVE SYSTEM

- The digestive system contributes to homeostasis by breaking down food into forms that can be absorbed and used by body cells. It also absorbs water, vitamins, and minerals, and it eliminates wastes from the body.
 - ❖ The food we eat consists of molecules that are too large to be used by body cells. Therefore, foods must be broken down into molecules that are small enough to enter body cells, a process known as digestion.
 - ❖ It extends from the mouth to the anus forms an extensive surface area in contact with the external environment, and is closely associated with the cardiovascular system. The gastrointestinal (GI) tract or alimentary canal is a continuous tube that extends from the mouth to the anus through the thoracic and abdominopelvic cavities.
- GI (alimentary canal)*
- Organs of the gastrointestinal tract include the mouth, most of the pharynx, esophagus, stomach, small intestine, and large intestine.

THE DIGESTIVE SYSTEM


- The accessory digestive organs include the teeth, tongue, salivary glands, liver, gallbladder, and pancreas.

FUNCTIONS OF THE DIGESTIVE SYSTEM

1. Ingestion: taking food into mouth.
2. Secretion: release of water, acid, buffers, and enzymes into lumen of GI tract.
3. Mixing and propulsion: churning and movement of food through GI tract.
4. Digestion: mechanical and chemical breakdown of food.
5. Absorption: passage of digested products from GI tract into blood and lymph.
6. Defecation: elimination of feces from GI tract.

Figure 24.2 Layers of the gastrointestinal tract. Variations in this basic plan may be seen in the esophagus (Figure 24.9), stomach (Figure 24.12), small intestine (Figure 24.19), and large intestine (Figure 24.24).

The four layers of the GI tract, from deep to superficial, are the **mucosa**, **submucosa**, **muscularis**, and **serosa**.



رح نرکز
علیم

NEURAL INNERVATION OF THE GI TRACT

- The gastrointestinal tract is regulated by an **intrinsic set of nerves known as the enteric nervous system** and by an **extrinsic set of nerves that are part of the autonomic nervous system.**

intrinsic Enteric nervous system

Extrinsic Autonomic nervous system






Figure 24.3 Organization of the enteric nervous system.

 The enteric nervous system consists of neurons arranged into the myenteric and submucosal plexuses.



<https://youtu.be/DG4fAdwdeS4?si=30fvYTYo4Ob7nAac>

easy explanation



Autonomic NS

- myenteric plexus
- motility (contraction)
- mechanical digestion

Enteric NS

- submucosal plexus
- secretion
- chemical digestion
- chemoreceptors

[mechanoreceptors] sensory receptors

ENTERIC NERVOUS SYSTEM

- ❖ The neurons of the ENS are arranged into two plexuses: **the myenteric plexus and submucosal plexus.**
- The myenteric plexus is located between the longitudinal and circular smooth muscle layers of the muscularis.
- The submucosal plexus is found within the submucosa.
- ❖ The plexuses of the ENS consist of **motor neurons, interneurons, and sensory neurons.**

Question 3 / 40

The submucosal plexus of the enteric nervous system is found:

1. Between the longitudinal and circular smooth muscle layers of the muscularis.
2. Within the circular smooth muscle layers of the muscularis.
3. Within the submucosa.
4. Within the serosa.

ENTERIC NERVOUS SYSTEM

- Because the motor neurons of the myenteric plexus supply the longitudinal and circular smooth muscle layers of the muscularis, this plexus mostly controls GI tract motility (movement), particularly the frequency and strength of contraction of the muscularis.
- The motor neurons of the submucosal plexus supply the secretory cells of the mucosal epithelium, controlling the secretions of the organs of the GI tract.
- ❖ The interneurons of the ENS interconnect the neurons of the myenteric and submucosal plexuses. The wall of the GI tract contains two major types of sensory receptors: (1) chemoreceptors, which respond to certain chemicals in the food present in the lumen, and (2) mechanoreceptors, such as stretch receptors, that are activated when food distends (stretches) the wall of a GI organ.

AUTONOMIC NERVOUS SYSTEM

↳ parasympathetic (rest and digest)

- ✓ Although the neurons of the ENS can function independently, they are subject to regulation by the neurons of the autonomic nervous system.
- ✓ **The vagus (X) nerves supply parasympathetic fibers to most parts of the GI tract, with the exception of the last half of the large intestine, which is supplied with parasympathetic fibers from the sacral spinal cord.** ★
- ✓ The parasympathetic nerves that supply the GI tract form neural connections with the ENS.

AUTONOMIC NERVOUS SYSTEM

- In general, stimulation of the ^{rest and digest} parasympathetic nerves that innervate the GI tract causes an increase in GI secretion and motility by increasing the activity of ENS neurons.
submucosal plexus myenteric plexus
- In general, the sympathetic nerves that supply the GI tract cause a decrease in GI secretion and motility by inhibiting the neurons of the ENS. Emotions such as anger, fear, and anxiety may slow digestion because they stimulate the sympathetic nerves that supply the GI tract.

MOUTH

Figure 24.5 Structures of the mouth (oral cavity).

 The mouth is formed by the cheeks, hard and soft palates, and tongue.





Figure 24.6 The three major salivary glands—¹parotid, ²sublingual, and ³submandibular. The submandibular glands, shown in the light micrograph (b), consist mostly of serous acini (serous fluid-secreting portions of gland) and a few mucous acini (mucus-secreting portions of gland); the parotid glands consist of serous acini only; and the sublingual glands consist of mostly mucous acini and a few serous acini.

● Saliva lubricates and dissolves foods and begins the chemical breakdown of carbohydrates and lipids.



TONGUE

☐ The tongue is an accessory digestive organ composed of skeletal muscle covered with mucous membrane.

☐ Together with its associated muscles, it forms the floor of the oral cavity.

☐ The extrinsic muscles of the tongue, which originate outside the tongue (attach to bones in the area) and insert into connective tissues in the tongue. *Anatomy ♦ bring insertion to the origin*

● The extrinsic muscles move the tongue from side to side and in and out to maneuver food for chewing, shape the food into a rounded mass, and force the food to the back of the mouth for swallowing. They also form the floor of the mouth and hold the tongue in position.


● The intrinsic muscles of the tongue originate in and insert into connective tissue within the tongue. They alter the shape and size of the tongue for speech and swallowing.

TEETH

Figure 24.7 A typical tooth and surrounding structures.

Teeth are anchored in sockets of the alveolar processes of the mandible and maxillae.

- A typical tooth has three major external regions: the crown, root, and neck.



Sagittal section of a mandibular (lower) molar



MECHANICAL AND CHEMICAL DIGESTION IN THE MOUTH

- Mechanical digestion in the mouth results from chewing, or mastication, in which food is manipulated by the tongue, ground by the teeth, and mixed with saliva.
 - As a result, the food is reduced to a soft, flexible, easily swallowed mass called a bolus.
 - Food molecules begin to dissolve in the water in saliva, an important activity because enzymes can react with food molecules in a liquid medium only.
- Two enzymes, salivary amylase and lingual lipase, contribute to chemical digestion in the mouth.
- Salivary amylase, which is secreted by the salivary glands, initiates the breakdown of starch. Dietary carbohydrates are either monosaccharide and disaccharide sugars or complex polysaccharides such as starches. Most of the carbohydrates we eat are starches, but only monosaccharides can be absorbed into the bloodstream. Thus, ingested disaccharides and starches must be broken down into monosaccharides.

↓

MECHANICAL AND CHEMICAL DIGESTION IN THE MOUTH

- Saliva also contains lingual lipase, which is secreted by lingual glands in the tongue. This enzyme becomes activated in the acidic environment of the stomach and thus starts to work after food is swallowed. It breaks down dietary triglycerides (fats and oils) into fatty acids and diglycerides. A diglyceride consists of a glycerol molecule that is attached to two fatty acids.

chemical digestion

enzymes

salivary amylase

↳ by salivary glands

Function:

- initiate breakdown of starch



monosaccharide

(in mouth inactive)
becomes active in
acidic environment

lingual lipase

↳ by lingual glands

Function:

- breaks down dietary
triglycerides (fats/oils)



Fatty acids / diglyceride

dietary carbohydrates

مقويات

monosaccharide ⇒ only can be absorbed

disaccharide (sugar)

polysaccharides (starch)

} ingested
to become
monosaccharide

7. What are the end products of carbohydrate digestion?

- A. chylomicrons
- B. amino acids
- C. free fatty acids
- D. monosaccharides

Answer is D: Monosaccharides or “simple” sugars.

TABLE 24.1

- ماضي لأغلب الفوق -

Summary of Digestive Activities in the Mouth

STRUCTURE	ACTIVITY	RESULT
Cheeks and lips	Keep <u>food between teeth</u> .	<u>Foods uniformly chewed during mastication.</u>
Salivary glands	Secrete <u>saliva</u> .	<u>Lining of mouth and pharynx moistened and lubricated.</u> Saliva softens, moistens, and dissolves food and cleanses mouth and teeth. Salivary amylase splits starch into smaller fragments (maltose, maltotriose, and α -dextrins).
Tongue		
Extrinsic tongue muscles	Move tongue <u>from side to side and in and out</u> .	Food <u>maneuvered for mastication</u> , shaped into <u>bolus</u> , and maneuvered for swallowing.
Intrinsic tongue muscles	<u>Alter shape of tongue</u> .	<u>Swallowing and speech.</u>
Taste buds	Serve as receptors for gustation (taste) and presence of food in mouth.	<u>Secretion of saliva</u> <u>stimulated by nerve impulses from taste buds to salivatory nuclei in brain stem to salivary glands.</u>
Lingual glands	Secrete <u>lingual lipase</u> .	<u>Triglycerides broken down into fatty acids and diglycerides.</u>
Teeth	Cut, tear, and <u>pulverize food</u> .	<u>Solid foods reduced to smaller particles for swallowing.</u>

سحق

PHARYNX

- When food is first swallowed, it passes from the mouth into the pharynx, a funnel-shaped tube that extends from the internal nares to the esophagus posteriorly and to the larynx anteriorly.

- The pharynx is composed of skeletal muscle and lined by mucous membrane, and is divided into three parts: the nasopharynx, the oropharynx, and the laryngopharynx.
Also tongue

<i>respiratory</i>	<i>respiratory</i>	<i>respiratory</i>
—	<i>digestion</i>	<i>digestion</i>

- The nasopharynx functions only in respiration, but both the oropharynx and laryngopharynx have digestive as well as respiratory functions.

- Swallowed food passes from the mouth into the oropharynx and laryngopharynx; the muscular contractions of these areas help propel food into the esophagus and then into the stomach.

ESOPHAGUS


No enzyme production
No absorption

Function :

- The esophagus secretes mucus and transports food into the stomach. It does not produce digestive enzymes, and it does not carry on absorption.

Figure 24.9 Histology of the esophagus. A higher-magnification view of nonkeratinized stratified squamous epithelium is shown in **Table 4.1F**.

● The esophagus secretes mucus and transports food to the stomach.



42. What is the function of the esophagus in digestion?

A. It is a site of mechanical digestion.

B. It transfers food from the mouth to the stomach.

C. The oesophagus secretes amylase to begin carbohydrate digestion.

D. the oesophagus secretes hydrochloric acid.


Answer is B: It is merely a conduit to transfer food from the mouth to the stomach

Figure 24.10 Deglutition (swallowing). During the pharyngeal stage (b) the tongue rises against the palate, the nasopharynx is closed off, the larynx rises, the epiglottis seals off the larynx, and the bolus is passed into the esophagus. During the esophageal stage (c), food moves through the esophagus into the stomach via peristalsis.

بلع

Key: Deglutition is a mechanism that moves food from the mouth into the stomach.

The tongue shapes the chewed, lubricated food (bolus) and moves it to the back of the mouth cavity.



(a) Position of structures during voluntary stage

(b) Pharyngeal stage of swallowing

Peristalsis, a progression of coordinated contractions and relaxations of the circular and longitudinal layers of the muscularis, pushes the bolus onward.

- The tongue rises against the palate and closes the nasopharynx.
 - The uvula and palate seal off the nasal cavity.
 - The epiglottis covers the larynx.
- Breathing is temporarily interrupted.

TABLE 24.2

Summary of Digestive Activities in the Pharynx and Esophagus

STRUCTURE	ACTIVITY	RESULT
Pharynx	<u>Pharyngeal stage of deglutition.</u>	<u>Moves bolus from oropharynx to laryngopharynx and into esophagus; closes air passageways.</u>
Esophagus	Relaxation of upper esophageal sphincter.	Permits entry of bolus from laryngopharynx into esophagus.
	<u>Esophageal stage of deglutition (peristalsis).</u>	<u>Pushes bolus down esophagus.</u>
	Relaxation of lower esophageal sphincter.	Permits entry of bolus into stomach.
	Secretion of mucus.	<u>Lubricates esophagus for smooth passage of bolus.</u>

بلع

⇒


معدة

يسيت

STOMACH

The stomach is a J-shaped enlargement of the GI tract directly inferior to the diaphragm in the abdomen.

● The stomach connects the esophagus to the duodenum, the first part of the small intestine.



(c) Esophageal stage of swallowing

STOMACH

mixing chamber
reservoir

- Because a meal can be eaten much more quickly than the intestines can digest and absorb it, one of the functions of the stomach is to serve as a mixing chamber and holding reservoir.
- ❖ At appropriate intervals after food is ingested, the stomach forces a small quantity of material into the first portion of the small intestine.
- ❖ The position and size of the stomach vary continually; the diaphragm pushes it inferiorly with each inhalation and pulls it superiorly with each exhalation. Empty, it is about the size of a large sausage, but it is the most distensible part of the GI tract and can accommodate a large quantity of food.
- In the stomach, digestion of starch and triglycerides continues, digestion of proteins begins, the semisolid bolus is converted to a liquid, and certain substances are absorbed.


قابل لتوسع

stomach

semisolid bolus ⇒ liquid

Figure 24.11 External and internal anatomy of the stomach.

The four regions of the stomach are the cardia, fundus, body, and pyloric part.



(a) Anterior view of regions of stomach

FUNCTIONS OF THE STOMACH

1. Mixes saliva, food, and gastric juice to form **chyme**.
2. Serves as reservoir for food before release into small intestine.
3. Secretes gastric juice, which contains **HCl** (kills bacteria and denatures proteins), **pepsin** (begins the digestion of proteins), **intrinsic factor** (aids absorption of vitamin B₁₂), and **gastric lipase** (aids digestion of triglycerides).
4. Secretes **gastrin** into blood.

The pylorus communicates with the duodenum of the small intestine via a smooth muscle sphincter called the pyloric sphincter.

The concave medial border of the stomach is called the lesser curvature; the convex lateral border is called the greater curvature.

STOMACH

- ❖ The stomach wall is composed of the same basic layers as the rest of the GI tract, with certain modifications. The surface of the mucosa is a layer of simple columnar epithelial cells called surface mucous cells.

↳ secrete alkaline mucus

- ❖ Parietal cells produce intrinsic factor (needed for absorption of vitamin B12) and hydrochloric acid.

۲
۱۹۳۵
۲ HCl

↳ triglycerides digestion

- ❖ The chief cells secrete pepsinogen and gastric lipase.

↳ inactive form → pepsin active form (protein digestion)

- ❖ The secretions of the mucous, parietal, and chief cells form gastric juice, which totals 2000–3000 mL per day.

Which type of stomach cells that can secrete pepsinogen:

1. Parietal cells.
2. G cells.
3. Chief cells.
4. Surface mucus cells.
5. Mucus neck cells.

11. Which of the following pairs of substances are **NOT** secreted by the stomach as part of “gastric juice”?

- A. hydrochloric acid and pepsinogen
- B. hormones and intrinsic factor
- C. nuclease and amylase
- D. mucus and gastrin

Answer is C: Nuclease and amylase are enzymes secreted by the pancreas

3. Which type of cell produces hydrochloric acid?

- A. Zymogenic cells
- B. Parietal cells
- C. Chief cells
- D. Enteroendocrine cells


Answer is B: The parietal cells of gastric glands produce hydrochloric acid. (Zymogenic cells are the same as Chief cells, they produce pepsinogen).

6-The cell which is responsible on absorption the b₁₂ (intrinsic factor)

Figure 24.12 Histology of the stomach.





Key: Gastric juice is the combined secretions of mucous cells, parietal cells, and chief cells.



(a) Three-dimensional view of layers of stomach

FIGURE 24.12 CONTINUES



(b) Sectional view of the stomach mucosa showing gastric glands and cell types

MECHANICAL AND CHEMICAL DIGESTION IN THE STOMACH

- ✓ Several minutes after food enters the stomach, waves of peristalsis pass over the stomach every 15 to 25 seconds.
- ✓ Few peristaltic waves are observed in the fundus, which primarily has a storage function.
↳ storage
- ✓ Instead, most waves begin at the body of the stomach and intensify as they reach the antrum.
- ✓ Each peristaltic wave moves gastric contents from the body of the stomach down into the antrum, a process known as propulsion. دفع
- ✓ The pyloric sphincter normally remains almost, but not completely, closed. Because most food particles in the stomach initially are too large to fit through the narrow pyloric sphincter, they are forced back into the body of the stomach, a process referred to as retropulsion. ارتجاع

MECHANICAL AND CHEMICAL DIGESTION IN THE STOMACH

- Another round of propulsion then occurs, moving the food particles back down into the antrum. If the food particles are still too large to pass through the pyloric sphincter, retropulsion occurs again as the particles are squeezed back into the body of the stomach. Then yet another round of propulsion occurs, and the cycle continues to repeat.
- The net result of these movements is that gastric contents are mixed with gastric juice, eventually becoming reduced to a soupy liquid called chyme. Once the food particles in chyme are small enough, they can pass through the pyloric sphincter, a phenomenon known as gastric emptying. Gastric emptying is a slow process: only about 3 mL of chyme moves through the pyloric sphincter at a time.

MECHANICAL AND CHEMICAL DIGESTION IN THE STOMACH

Foods may remain in the fundus for about an hour without becoming mixed with gastric juice. During this time, digestion by salivary amylase from the salivary glands continues. Soon, however, the churning action mixes chyme with acidic gastric juice, inactivating salivary amylase and activating lingual lipase.


lingual lipase	salivary glands
gastric lipase	stomach
pancreatic lipase	pancreas

ghena is ☺
your uncle
Love u

Figure 24.13 Secretion of HCl (hydrochloric acid) by parietal cells in the stomach.



Proton pumps, powered by ATP, secrete the H^+ ; Cl^- diffuses into the stomach lumen through Cl^- channels.



Key:



Proton pump (H^+-K^+ ATPase)




K^+ (potassium ion) channel




Cl^- (chloride ion) channel



Carbonic anhydrase




Diffusion



HCO_3^- - Cl^- antiporter

Figure 24.14 Regulation of HCl secretion.

HCl secretion by parietal cells can be stimulated by several sources: **acetylcholine (ACh)**, **gastrin**, and **histamine**.



REGULATION OF HCL SECRETION

اجی
سنوات

HCl secretion by parietal cells can be stimulated by several sources:

1. Acetylcholine (ACh) is released by parasympathetic neurons.
2. Gastrin secreted by G cells.  stomach
3. Histamine, which is a paracrine substance released by mast cells in the nearby lamina propria.

❖ Acetylcholine and gastrin stimulate parietal cells to secrete more HCl in the presence of histamine. In other words, histamine acts synergistically, enhancing the effects of acetylcholine and gastrin. Receptors for all three substances are present in the plasma membrane of parietal cells.

بیشکل
تکاملی

5. What is the role of gastrin in the digestive system?
- A. to stimulate release of bile and pancreatic juice
 - B. to stimulate parietal cells that secrete HCL
 - C. to activate pepsinogen
 - D. to hydrolyse proteins to polypeptides

B

REGULATION OF HCL SECRETION

- The strongly acidic fluid of the stomach kills many microbes in food. Function 8
- HCl partially denatures (unfolds) proteins in food and stimulates the secretion of hormones that promote the flow of bile and pancreatic juice.
- ☆ ➤ Enzymatic digestion of proteins also begins in the stomach.
- The only proteolytic (protein-digesting) enzyme in the stomach is pepsin, which is secreted by chief cells.
- Pepsin ^{يقطع} severs certain peptide bonds between amino acids, breaking down a protein chain of many amino acids into smaller peptide fragments.
- Pepsin is most effective in the very acidic environment of the stomach (pH 2); it becomes inactive at a higher pH.

↓
pepsinogen

WHAT KEEPS PEPSIN FROM DIGESTING THE PROTEIN IN STOMACH CELLS ALONG WITH THE FOOD?

- ❖ First, pepsin is secreted in an inactive form called pepsinogen; in this form, it cannot digest the proteins in the chief cells that produce it.
- ❖ Pepsinogen is not converted into active pepsin until it comes in contact with hydrochloric acid secreted by parietal cells or active pepsin molecules.
HCl acid
pH=2
- ❖ Second, the stomach epithelial cells are protected from gastric juices by a layer 1–3 mm thick of alkaline mucus secreted by surface mucous cells and mucous neck cells.

41. Which of the following could **NOT** be used to describe pepsinogen?

- A. it is a protein
- B. it is a hormone
- C. it is related to an enzyme
- D. it is inactive

Answer is B: Pepsinogen is an inactive protein enzyme. But it is not a hormone.

55. Correctly complete the sentence: Pepsinogen is
- A. converted to pepsin by hydrochloric acid
 - B. converted to pepsin by intrinsic factor
 - C. secreted by the pancreas
 - D. involved in production of carbohydrate digesting enzymes

A

REGULATION OF HCL SECRETION

- Another enzyme of the stomach is gastric lipase, which splits triglycerides (fats and oils) in fat molecules (such as those found in milk) into fatty acids and monoglycerides.

gastric lipase \Rightarrow triglycerides \rightarrow Fats \rightarrow fatty acids / monoglycerid

- This enzyme, which has a limited role in the adult stomach, operates best at a pH of 5–6. More important than either lingual lipase or gastric lipase is pancreatic lipase, an enzyme secreted by the pancreas into the small intestine.

- Within 2 to 4 hours after eating a meal, the stomach has emptied its contents into the duodenum. Foods rich in carbohydrates spend the least time in the stomach; high-protein foods remain somewhat longer, and emptying is slowest after a fat-laden meal containing large amounts of triglycerides.

TABLE 24.3

Summary of Digestive Activities in the Stomach

STRUCTURE	ACTIVITY	RESULT
Mucosa (stomach)		
Surface mucous cells and mucous neck cells	1- <u>Secrete mucus.</u> 2- <u>Absorption.</u>	Forms protective barrier that prevents digestion of stomach wall. Small quantity of water, ions, short-chain fatty acids, and some drugs enter bloodstream.
Parietal cells	1- Secrete <u>intrinsic factor.</u>	Needed for <u>absorption of vitamin B₁₂</u> (used in red blood cell formation, or erythropoiesis).
Chief cells	2- Secrete <u>hydrochloric acid.</u> HCl Secrete <u>pepsinogen.</u>	1- <u>Kills microbes in food; denatures proteins; converts pepsinogen into pepsin.</u> HCl 2- <u>Pepsin (activated form) breaks down proteins into peptides.</u>
G cells	Secrete <u>gastric lipase.</u> Secrete <u>gastrin.</u>	<u>Splits triglycerides into fatty acids and monoglycerides.</u> Stimulates <u>parietal cells to secrete HCl</u> and <u>chief cells to secrete pepsinogen</u> ; contracts lower esophageal sphincter, increases motility of stomach, and relaxes pyloric sphincter.
G	Gastrin	
Muscularis	Mixing waves (<u>gentle peristaltic movements</u>).	Churns and physically breaks down food and mixes it with gastric juice, forming chyme. Forces chyme through pyloric sphincter.
Pyloric sphincter	Opens to permit <u>passage of chyme into duodenum.</u>	Regulates passage of chyme from stomach to duodenum; prevents backflow of chyme from duodenum to stomach.

93. Which fluid within the body is likely to have the **LOWEST** pH?

- A. the chyme in the ileum.
- B. saliva.
- C. the blood.
- D. the chyme in the stomach.

Answer is D: The stomach contents are the most acidic, that is, has the lowest pH.

سؤال true false

هيك مضمونه

The lowest PH in the small intestine

جوابه false


PANCREAS

- From the stomach, chyme passes into the small intestine. Because chemical digestion in the small intestine depends on activities of the pancreas, liver, and gallbladder, we first consider the activities of these accessory digestive organs and their contributions to digestion in the small intestine.

Figure 24.15 Relationship of the pancreas to the liver, gallbladder, and duodenum. The inset (b) shows details of the common bile duct and pancreatic duct forming the hepatopancreatic ampulla and emptying into the duodenum.




 Pancreatic enzymes digest starches (polysaccharides), proteins, triglycerides, and nucleic acids.




PANCREAS

- The pancreas, a **retroperitoneal gland** that is about 12–15 cm (5–6 in.) long and 2.5 cm (1 in.) thick, lies posterior to the greater curvature of the stomach.
- The pancreas consists of a head, a body, and a tail and is usually connected to the duodenum by two ducts.
- The head is the expanded portion of the organ near the curve of the duodenum; superior to and to the left of the head are the central body and the tapering tail.

لنو صلیح





(c) Ducts carrying bile from liver and gallbladder and pancreatic juice from pancreas to the duodenum

PANCREAS

- The protein-digesting enzymes of the pancreas are produced in an inactive form just as pepsin is produced in the stomach as pepsinogen. Because they are inactive, the enzymes do not digest cells of the pancreas itself.

● Trypsin is secreted in an inactive form called trypsinogen.

Active

Inactive

- Pancreatic acinar cells also secrete a protein called trypsin inhibitor that combines with any trypsin formed accidentally in the pancreas or in pancreatic juice and blocks its enzymatic activity.

Active

● When trypsinogen reaches the lumen of the small intestine, it encounters an activating brush-border enzyme called enterokinase, which splits off part of the trypsinogen molecule to form trypsin.

trypsinogen meet enterokinase = Trypsin (Active)

76. What are the cells in the pancreas that secrete “pancreatic juice” called?

A. hepatocytes

B. Peyer’s patches

C. the acini


D. islets of Langerhans

LIVER AND GALLBLADDER


- The liver is the heaviest gland of the body, weighing about 1.4 kg (about 3 lb) in an average adult. Of all of the organs of the body, it is second only to the skin in size.
- The liver is inferior to the diaphragm and occupies most of the right hypochondriac and part of the epigastric regions of the abdominopelvic cavity.
- The gallbladder is a pear-shaped sac that is located in a depression of the posterior surface of the liver. It is 7–10 cm (3–4 in.) long and typically hangs from the anterior inferior margin of the liver.

Figure 24.16 Histology of the liver.

Histologically, the liver is composed of hepatocytes, bile canaliculi, and hepatic sinusoids.




(a) Overview of histological components of liver



(b) Details of histological components of liver

Figure 24.17 Hepatic blood flow: sources, path through the liver, and return to the heart.

The liver receives oxygenated blood via the hepatic artery and nutrient-rich deoxygenated blood via the hepatic portal vein.



LIVER AND GALLBLADDER

- From the **hepatic artery** it obtains oxygenated blood, and from the **hepatic portal vein** it receives deoxygenated blood containing newly absorbed nutrients, drugs, and possibly microbes and toxins from the gastrointestinal tract.
- **Branches of both the hepatic artery and the hepatic portal vein carry blood into hepatic sinusoids, where oxygen, most of the nutrients, and certain toxic substances are taken up by the hepatocytes.**
- Products manufactured by the hepatocytes and nutrients needed by other cells are secreted back into the blood, which then drains into the central vein and eventually passes into a hepatic vein.
- Because blood from the gastrointestinal tract passes through the liver as part of the hepatic portal circulation, the liver is often a site for metastasis of cancer that originates in the GI tract.

40. A lobule of the liver contains several blood vessels. Which one carries nutrient-rich blood from the small intestine?

A. hepatic artery proper

B. hepatic portal vein

C. central vein

D. bile ductule

Branches of both the hepatic artery and the hepatic portal vein carry blood into hepatic sinusoids, where oxygen, most of the nutrients, and certain toxic substances are taken up by the hepatocytes.

1. True.
2. False.

Next

FUNCTIONS OF THE LIVER AND GALLBLADDER

- Each day, hepatocytes secrete 800–1000 mL (about 1 qt) of bile, a yellow, brownish, or olive-green liquid. It has a pH of 7.6–8.6 and consists mostly of water, bile salts, cholesterol, a phospholipid called lecithin, bile pigments, and several ions.
- The principal bile pigment is bilirubin. 1 2 3
يحرر
- The phagocytosis of aged red blood cells liberates iron, globin, and bilirubin (derived from heme).
- The iron and globin are recycled; the bilirubin is secreted into the bile and is eventually broken down in the intestine. One of its breakdown products—stercobilin—gives feces their normal brown color.
- Bile is partially an excretory product and partially a digestive secretion.
- Bile salts, which are sodium salts and potassium salts of bile acids play a role in emulsification, the breakdown of large lipid globules into a suspension of small lipid globules.
→ process of forming a stable mixture of liquids
استحلاب
- The small lipid globules present a very large surface area that allows pancreatic lipase to more rapidly accomplish digestion of triglycerides. Bile salts also aid in the absorption of lipids following their digestion.

33. What is the function of bile salts?
- A. To assist the absorption of digested lipids
 - B. To emulsify lipids
 - C. To hydrolyse lipids
 - D. To digest lipids

B

6. Which liver cells produce bile?

A. Kupffer cells

B. sinusoids

C. hepatocytes

D. the acini

Answer is C: Hepatocytes are liver cells. Kupffer cells are macrophages, while sinusoids are blood capillaries.

78. Which of the following is a function of bile?
- A. To attach to vitamin B12 to allow it to be absorbed.
 - B. To activate trypsinogen, chymotrypsinogen and procarboxypeptidase.
 - C. To digest fats.
 - D. To disperse large lipid globules into smaller droplets.


تَشْتِيت

Answer is D: This is known as emulsification.

5- function the bile (degradation the lipid)

FUNCTIONS OF THE LIVER AND GALLBLADDER

- Digestion and absorption continue in the small intestine, **bile release increases.**
- Between meals, after most absorption has occurred, **bile flows into the gallbladder for storage** because the sphincter of the hepatopancreatic ampulla closes off the entrance to the duodenum. The sphincter surrounds the hepatopancreatic ampulla.



23. Which of the following is TRUE of bile?

A. it converts inactive pancreatic enzymes to active form.

B. needed in the small intestine for the digestion of fats.

C. synthesised by the gall bladder.

D. needed in the small intestine for the emulsification of fats.

Answer is D: bile emulsifies (rather than digests) fat. It is stored (but not synthesised) in the gall bladder.

Liver Functions

Carb metabolism

glucose ↓ glycogen → glucose
glucose ↑ glucose → glycogen
↳ stored in liver

lipid metabolism

- hepatocytes store some Triglyceride
- synthesize cholesterol
- use cholesterol to make bile salts

protein metabolism

- hepatocytes deaminate amino acid
(remove ↓ NH_2)
- the toxic ammonia NH_3 is converted to urea
- hepatocytes synthesize plasma protein
Alpha / beta / globulins
Albumin / prothrombin / Fibrinogen

processing of drugs and hormones

(Detoxification)

- detoxify substances: Alcohol
- Excrete drugs
penicillin / erythromycin
sulfonamides into bile

excretion of bilirubin

- bilirubin in bile is metabolised then secreted
↳ by small intestine

synthesis of bile salts

- used in small intestine
- function emulsification and absorption of lipids

- Excrete thyroid hormones and steroid hormones :
estrogen / Aldosterone

Storage

- store glycogen
- store Vitamins

(AKED, B12)
أكيد

- store minerals
(iron, copper)

phagocytosis

- Kupfer cells

(stellate (reticuloendothelial))

phagocytize Aged

red blood cells

white blood cells

some bacteria

Activation of vitamin D

skin

liver

Kidneys

synthesize the active

Form of vitamin D

● FUNCTIONS OF THE LIVER AND GALLBLADDER

- In addition to secreting bile, which is needed for absorption of dietary fats, the liver performs many other vital functions.
1. **Carbohydrate metabolism:** The liver is especially important in maintaining a normal blood glucose level. When blood glucose is low, the liver can break down glycogen to glucose and release the glucose into the bloodstream. When blood glucose is high, as occurs just after eating a meal, the liver converts glucose to glycogen and triglycerides for storage.
 2. **Lipid metabolism:** Hepatocytes store some triglycerides, synthesize cholesterol; and use cholesterol to make bile salts.
 3. **Protein metabolism:** Hepatocytes deaminate (remove the amino group, NH₂, from) amino acids, the resulting toxic ammonia (NH₃) is then converted into the much less toxic urea, which is excreted in urine. Hepatocytes also synthesize most plasma proteins, such as alpha and beta globulins, albumin, prothrombin, and fibrinogen.

48. The liver is able to deaminate amino acids forming ammonia in the process. What happens to the ammonia?
- A. It is phagocytosed by Kupffer cells.
 - B. It is used in transamination to form non-essential amino acids.
 - C. It is converted to bile to be excreted via the gut.
 - D. It is converted to urea for excretion by the kidneys.

Answer is D: Urea is the molecule and vehicle for excretion of human nitrogenous waste.

50. If blood glucose is high, what does the liver do about it?
- A. The liver converts glucose to glycogen or triglycerides.
 - B. The liver performs glycogenolysis.
 - C. The liver performs gluconeogenesis.
 - D. The liver transaminates glucose to produce amino acids.

A

● FUNCTIONS OF THE LIVER AND GALLBLADDER

- In addition to secreting bile, which is needed for absorption of dietary fats, the liver performs many other vital functions.
- 4. Processing of drugs and hormones: The liver can detoxify substances such as alcohol and excrete drugs such as penicillin, erythromycin, and sulfonamides into bile. It can also chemically alter or excrete thyroid hormones and steroid hormones such as estrogens and aldosterone.
- 5. Excretion of bilirubin: As previously noted, bilirubin, derived from the heme of aged red blood cells, is absorbed by the liver. Most of the bilirubin in bile is metabolized in the small intestine by bacteria and eliminated in feces.
- 6. Synthesis of bile salts: Bile salts are used in the small intestine for the emulsification and absorption of lipids.

● FUNCTIONS OF THE LIVER AND GALLBLADDER

- In addition to secreting bile, which is needed for absorption of dietary fats, the liver performs many other vital functions.

اكد
(AKED, B12)


7. **Storage:** In addition to glycogen, the liver is a prime storage site for certain vitamins (A, B12, D, E, and K) and minerals (iron and copper), which are released from the liver when needed elsewhere in the body.
8. **Phagocytosis:** The stellate reticuloendothelial (Kupffer) cells of the liver phagocytize aged red blood cells, white blood cells, and some bacteria.
9. **Activation of vitamin D:** The skin, liver, and kidneys participate in synthesizing the active form of vitamin D.

SMALL INTESTINE

- Most digestion and absorption of nutrients occur in a long tube called the small intestine.
- Its length alone provides a large surface area for digestion and absorption, and that area is further increased by circular folds, villi, and microvilli.

Figure 24.18 Anatomy of the small intestine. (a) Regions of the small intestine are the duodenum, jejunum, and ileum. (b) Circular folds increase the surface area for digestion and absorption in the small intestine.


 Most digestion and absorption occur in the small intestine.



(a) Anterior view of external anatomy

FUNCTIONS OF THE SMALL INTESTINE

1. Segmentations mix chyme with digestive juices and bring food into contact with mucosa for absorption; peristalsis propels chyme through small intestine.
2. Completes digestion of carbohydrates, proteins, and lipids; begins and completes digestion of nucleic acids.
3. Absorbs about 90% of nutrients and water that pass through digestive system.



(b) Internal anatomy of jejunum

Figure 24.19 Histology of the small intestine.

Circular folds, villi, and microvilli increase the surface area of the small intestine for digestion and absorption.





Figure 24.20 Histology of the duodenum and ileum.

 Microvilli in the small intestine contain several brush-border enzymes that help digest nutrients.



(a) Wall of the duodenum

8. What feature of the small intestine enhances its ability to absorb digested food?

- A. its large surface area
- B. the gaps between adjacent epithelial cells
- C. secretion of the hormone absorptin
- D. its longer length compared to the large intestine

A

ROLE OF INTESTINAL JUICE AND BRUSH-BORDER ENZYMES

- About 1–2 liters of intestinal juice, a clear yellow fluid, is secreted each day. Intestinal juice contains water and mucus and is slightly alkaline (pH 7.6).



- ★ ● The alkaline pH of intestinal juice is due to its high concentration of bicarbonate ions.



- The absorptive cells of the small intestine synthesize several digestive enzymes, called brush-border enzymes, and insert them in the plasma membrane of the microvilli.

ROLE OF INTESTINAL JUICE AND BRUSH-BORDER ENZYMES

- Thus, some enzymatic digestion occurs at the surface of the absorptive cells that line the villi, rather than in the lumen exclusively, as occurs in other parts of the GI tract. Among the brush-border enzymes are four carbohydrate-digesting enzymes called α -dextrinase, maltase, sucrase, and lactase; protein-digesting enzymes called peptidases (aminopeptidase and dipeptidase); and two types of nucleotide-digesting enzymes, nucleosidases and phosphatases.

brush border enzymes

digesting -enzymes

4 carbohydrates enzymes

- α dextrinase
- maltase
- sucrase
- lactase

protein enzymes

- (peptides)
- aminopeptidases
- dipeptidases

nucleotide enzymes

- nucleosidases
- phosphatases

MECHANICAL DIGESTION IN THE SMALL INTESTINE

- The two types of movements of the small intestine—segmentations and a type of peristalsis called migrating motility complexes—are governed mainly by the myenteric plexus.

motility

- Segmentations are localized, mixing contractions that occur in portions of intestine distended by a large volume of chyme. Segmentations mix chyme with the digestive juices and bring the particles of food into contact with the mucosa for absorption; they do not push the intestinal contents along the tract.

1- Movement the digestive food with
mixing the enzyme (segmentation)

MECHANICAL DIGESTION IN THE SMALL INTESTINE

- Segmentations occur most rapidly in the duodenum, about 12 times per minute, and progressively slow to about 8 times per minute in the ileum.
- After most of a meal has been absorbed, which lessens distension of the wall of the small intestine, segmentation stops and peristalsis begins. الأنتفاخ
- The type of peristalsis that occurs in the small intestine, termed a migrating motility complex (MMC), begins in the lower portion of the stomach and pushes chyme forward along a short stretch of small intestine before dying out.
- The MMC slowly migrates down the small intestine, reaching the end of the ileum in 90–120 minutes. Then another MMC begins in the stomach. Altogether, chyme remains in the small intestine for 3–5 hours.

CHEMICAL DIGESTION IN THE SMALL INTESTINE

1. In the mouth, **salivary amylase converts starch (a polysaccharide)** to maltose (a disaccharide), maltotriose (a trisaccharide), and α -dextrins (short-chain, branched fragments of starch with 5–10 glucose units).
2. In the stomach, **pepsin converts proteins to peptides** (small fragments of proteins), and **lingual and gastric lipases convert some triglycerides** into fatty acids, diglycerides, and monoglycerides.
3. Thus, chyme entering the small intestine contains partially digested carbohydrates, proteins, and lipids.
4. The completion of the digestion of carbohydrates, proteins, and lipids is a collective effort of pancreatic juice, bile, and intestinal juice in the small intestine.

ABSORPTION IN THE SMALL INTESTINE

- ❑ All of the chemical and mechanical phases of digestion from the mouth through the small intestine are directed toward changing food into forms that can pass through the absorptive epithelial cells lining the mucosa and into the underlying blood and lymphatic vessels.
- ❑ These forms are monosaccharides (glucose, fructose, and galactose) from carbohydrates; single amino acids, dipeptides, and tripeptides from proteins; and fatty acids, glycerol, and monoglycerides from triglycerides.
- ❑ Passage of these digested nutrients from the gastrointestinal tract into the blood or lymph is called absorption.

ABSORPTION IN THE SMALL INTESTINE


- Absorption of materials occurs via diffusion, facilitated diffusion, osmosis, and active transport.
- ★ ● About 90% of all absorption of nutrients occurs in the small intestine; the other 10% occurs in the stomach and large intestine.
- Any undigested or unabsorbed material left in the small intestine passes on to the large intestine.

11-Most absorption occurs in the
stomach(false)


Figure 24.21 Absorption of digested nutrients in the small intestine. For simplicity, all digested foods are shown in the lumen of the small intestine, even though some nutrients are digested by brush-border enzymes.



Key: Long-chain fatty acids and monoglycerides are absorbed into lacteals; other products of digestion enter blood capillaries.



(a) Mechanisms for movement of nutrients through absorptive epithelial cells of villi



(b) Movement of absorbed nutrients into blood and lymph

Figure 24.22 Daily volumes of fluid ingested, secreted, absorbed, and excreted from the GI tract.

 All water absorption in the GI tract occurs via osmosis.




TABLE 24.4

Summary of Digestive Activities in the Pancreas, Liver, Gallbladder, and Small Intestine

STRUCTURE	ACTIVITY
Pancreas	Delivers <u>pancreatic juice</u> into <u>duodenum</u> via <u>pancreatic duct</u> to assist absorption (see Table 24.5 for pancreatic enzymes and their functions).
Liver	Produces <u>bile (bile salts)</u> necessary for <u>emulsification</u> and absorption of lipids.
Gallbladder	Stores, concentrates, and delivers <u>bile</u> into <u>duodenum</u> via <u>common bile duct</u> .
Small Intestine	90% <u>Major site of digestion and absorption of nutrients and water in gastrointestinal tract.</u>
Mucosa/submucosa	
Intestinal glands	Secrete <u>intestinal juice</u> to assist absorption.
Absorptive cells	Digest and absorb nutrients. / <i>synthesize brush border enzymes</i>
Goblet cells	Secrete <u>mucus</u> .
Enteroendocrine cells (S, CCK, K)	Secrete secretin, cholecystokinin, and glucose-dependent insulinotropic peptide.
Paneth cells	Secrete lysozyme (bactericidal enzyme) and phagocytosis.
Duodenal (Brunner's) glands	Secrete alkaline fluid to buffer stomach acids, and mucus for protection and lubrication.
Circular folds	Folds of mucosa and submucosa that <u>increase surface area</u> for digestion and absorption.
Villi	Fingerlike projections of mucosa that are sites of absorption of digested food and <u>increase surface area</u> for digestion and absorption.
Microvilli	Microscopic, membrane-covered projections of absorptive epithelial cells that <u>contain brush-border enzymes</u> (listed in Table 24.5) and that <u>increase surface area</u> for digestion and absorption.
Muscularis	
Segmentation	<u>Type of peristalsis</u> : alternating <u>contractions of circular smooth muscle fibers</u> that produce segmentation and resegmentation of sections of small intestine; <u>mixes chyme with digestive juices</u> and brings food into contact with mucosa for absorption.
Migrating motility complex (MMC)	<u>Type of peristalsis</u> : waves of <u>contraction and relaxation of circular and longitudinal smooth muscle</u> fibers passing the length of the small intestine; <u>moves chyme toward ileocecal sphincter</u> .

TABLE 24.5

Summary of Digestive Enzymes

ENZYME	SOURCE	SUBSTRATES	PRODUCTS
SALIVA			
Salivary amylase	Salivary glands. <i>mouth</i>	<u>Starches (polysaccharides).</u>	- Maltose (disaccharide), - maltotriose (trisaccharide), and - α -dextrins.
Lingual lipase	Lingual glands in <u>tongue</u> .	<u>Triglycerides (fats and oils)</u> and other lipids.	<u>Fatty acids and diglycerides.</u>
GASTRIC JUICE			
Pepsin (activated from pepsinogen by pepsin and hydrochloric acid)	<u>Stomach chief cells.</u>	Proteins.	Peptides.
Gastric lipase	<u>Stomach chief cells.</u>	<u>Triglycerides (fats and oils).</u>	<u>Fatty acids and monoglycerides.</u>
PANCREATIC JUICE			
Pancreatic amylase	Pancreatic acinar cells.	<u>Starches (polysaccharides).</u>	Maltose (disaccharide), maltotriose (trisaccharide), and α -dextrins.
Trypsin (activated from trypsinogen by enterokinase)	Pancreatic acinar cells.	Proteins.	Peptides.
Chymotrypsin (activated from chymotrypsinogen by trypsin)	Pancreatic acinar cells.	Proteins.	Peptides.
Elastase (activated from proelastase by trypsin)	Pancreatic acinar cells.	Proteins.	Peptides.
Carboxypeptidase (activated from procarboxypeptidase by trypsin)	Pancreatic acinar cells.	Amino acid at carboxyl end of peptides.	Amino acids and peptides.
Pancreatic lipase	Pancreatic acinar cells.	Triglycerides (fats and oils) that have been emulsified by bile salts.	<u>Fatty acids and monoglycerides.</u>
Nucleases			
Ribonuclease	Pancreatic acinar cells.	Ribonucleic acid.	Nucleotides.
Deoxyribonuclease	Pancreatic acinar cells.	Deoxyribonucleic acid.	Nucleotides.

proteins
↓
peptides

TABLE 24.5**Summary of Digestive Enzymes****Nucleases**

Ribonuclease

Pancreatic acinar cells.

Ribonucleic acid.

Nucleotides.

Deoxyribonuclease

Pancreatic acinar cells.

Deoxyribonucleic acid.

Nucleotides.

BRUSH-BORDER ENZYMES IN MICROVILLI PLASMA MEMBRANE α -Dextrinase

Small intestine.

 α -Dextrins.

Glucose.

Maltase

Small intestine.

Maltose.

Glucose.

Sucrase

Small intestine.

Sucrose.

Glucose and fructose.

Lactase

Small intestine.

Lactose.

Glucose and galactose.

Enterokinase

Small intestine.

Trypsinogen.

Trypsin.

Peptidases

Aminopeptidase

Small intestine.

Amino acid at amino end of peptides.

Amino acids and peptides.

Dipeptidase

Small intestine.

Dipeptides.

Amino acids.

Nucleosidases and
phosphatases

Small intestine.

Nucleotides.

Nitrogenous bases, pentoses, and
phosphates.

LARGE INTESTINE

No enzymes


- The large intestine is **the terminal portion of the GI tract.**
- The overall functions of the large intestine are the completion of absorption, the production of certain vitamins, the formation of feces, and the expulsion of feces from the body.

Figure 24.23 Anatomy of the large intestine.

The regions of the large intestine are the cecum, colon, rectum, and anal canal.

FUNCTIONS OF THE LARGE INTESTINE

1. Haustral churning, peristalsis, and mass peristalsis drive contents of colon into rectum.
2. Bacteria in large intestine convert proteins to amino acids, break down amino acids, and produce some B vitamins and vitamin K. - KB -
3. Absorption of some water, ions, and vitamins.
4. Formation of feces.
5. Defecation (emptying rectum).




(a) Anterior view of large intestine showing


(b) Frontal section of anal canal

Figure 24.24 Histology of the large intestine.

Intestinal glands formed by simple columnar epithelial cells and goblet cells extend the full thickness of the mucosa.



(a) Three-dimensional view of layers of large intestine



(b) Sectional view of intestinal glands and cell types


MECHANICAL DIGESTION IN THE LARGE INTESTINE

- ❑ The passage of chyme from the ileum into the cecum is regulated by the action of the ileocecal sphincter.
- ❑ Normally, the valve remains partially closed **so that the passage of chyme into the cecum usually occurs slowly.**
- ❑ Immediately after a meal, **a gastroileal reflex intensifies** peristalsis in the ileum and forces **any chyme into the cecum.**
- ★ ❑ **The hormone gastrin also relaxes the sphincter.**
- ❑ Whenever the **cecum is distended**, **the degree of contraction** of the ileocecal sphincter **intensifies.**

MECHANICAL DIGESTION IN THE LARGE INTESTINE

- ❑ Movements of the colon begin when substances pass the ileocecal sphincter.
- ❑ Because chyme moves through the small intestine at a fairly constant rate, the time required for a meal to pass into the colon is determined by gastric emptying time.
- As food passes through the ileocecal sphincter, it fills the cecum and accumulates in the ascending colon.
- One movement characteristic of the large intestine is haustral churning. In this process, the haustra remain relaxed and become distended while they fill up. When the distension reaches a certain point, the walls contract and squeeze the contents into the next haustrum.
- Peristalsis also occurs, although at a slower rate (3–12 contractions per minute) than in more proximal portions of the tract.

Haustral churning



- Each segment in large intestine is called a haustra
- Smooth muscle contractions move contents along (type of peristalsis)
- Water is absorbed in large intestine

MECHANICAL DIGESTION IN THE LARGE INTESTINE

- A final type of movement is **mass peristalsis**, a strong peristaltic wave that **begins at about the middle of the transverse colon** and quickly **drives the contents of the colon into the rectum.**
- Because food in the stomach initiates this gastrocolic reflex in the colon, **mass peristalsis usually takes place three or four times a day, during or immediately after a meal.**

CHEMICAL DIGESTION IN THE LARGE INTESTINE

- The final stage of digestion occurs in the colon through the activity of bacteria that inhabit the lumen. Mucus is secreted by the glands of the large intestine, but no enzymes are secreted.
- Chyme is prepared for elimination by the action of bacteria, which ferment any remaining carbohydrates and release hydrogen, carbon dioxide, and methane gases. These gases contribute to flatus (gas) in the colon, termed flatulence when it is excessive.
- Bacteria also convert any remaining proteins to amino acids and break down the amino acids into simpler substances: indole, skatole, hydrogen sulfide, and fatty acids.
- * ● Bacteria also decompose bilirubin to simpler pigments, including stercobilin, which gives feces their brown color.
- Bacterial products that are absorbed in the colon include several vitamins needed for normal metabolism, among them some B vitamins and vitamin K.

انتفاخ البطن

KB ك ب

ABSORPTION AND FECES FORMATION IN THE LARGE INTESTINE

- ❑ By the time **chyme has remained in the large intestine 3–10 hours, it has become solid or semisolid because of water absorption and is now called feces.**
- ❑ Chemically, feces consist of water, inorganic salts, sloughed-off epithelial cells from the mucosa of the gastrointestinal tract, bacteria, products of bacterial decomposition, unabsorbed digested materials, and indigestible parts of food.
- ❑ Although 90% of all water absorption occurs in the small intestine, **the large intestine absorbs enough to make it an important organ in maintaining the body's water balance.**
- ❑ **The large intestine also absorbs ions, including sodium and chloride, and some vitamins.**



THE DEFECATION REFLEX

- ❑ Mass peristaltic movements push fecal material from the sigmoid colon into the rectum.

انتفاخ

- ❑ The resulting distension of the rectal wall stimulates stretch receptors, which initiates a defecation reflex that results in defecation, the elimination of feces from the rectum through the anus.

THE DEFECATION REFLEX

- ❖ The amount of bowel movements that a person has over a given period of time **depends on various factors such as diet, health, and stress.**
- ❖ The normal range of bowel activity varies from two or three bowel movements per day to three or four bowel movements per week.
- **Diarrhea** is an increase in the frequency, volume, and fluid content of the feces caused by increased motility of and decreased absorption by the intestines.
- **Constipation** refers to infrequent or difficult defecation caused by decreased motility of the intestines.

TABLE 24.6

Summary of Digestive Activities in the Large Intestine

STRUCTURE	ACTIVITY	FUNCTION(S)
Lumen	<u>Bacterial activity.</u>	Breaks down undigested carbohydrates, proteins, and amino acids into products that can be expelled in feces or absorbed and detoxified by liver; synthesizes certain B vitamins and vitamin K.
Mucosa	<u>Secretes mucus.</u> <u>Absorption.</u>	Lubricates colon; protects mucosa. Water absorption solidifies feces and contributes to body's water balance; <u>solute absorbed include ions and some vitamins.</u>
Muscularis	1 <u>Haustral churning.</u> 2 <u>Peristalsis.</u> 3 <u>Mass peristalsis.</u> 4 <u>Defecation reflex.</u>	Moves contents from <u>hastrum to hastrum</u> by <u>muscular contractions.</u> Moves contents along length of <u>colon</u> by <u>contractions of circular and longitudinal muscles.</u> <u>Forces contents into sigmoid colon and rectum.</u> <u>Eliminates feces by contractions in sigmoid colon and rectum.</u>

TABLE 24.7

Summary of Organs of the Digestive System and Their Functions

ORGAN	FUNCTION(S)
Tongue	Maneuvers food for mastication, shapes food into a bolus, maneuvers food for deglutition, detects sensations for taste, and initiates digestion of triglycerides.
Salivary glands	Saliva produced by these glands softens, moistens, and dissolves foods; cleanses mouth and teeth; initiates the digestion of starch.
Teeth	Cut, tear, and pulverize food to reduce solids to smaller particles for swallowing.
Pancreas	Pancreatic juice buffers acidic gastric juice in chyme, stops the action of pepsin from the stomach, creates the proper pH for digestion in the small intestine, and participates in the digestion of carbohydrates, proteins, triglycerides, and nucleic acids.
Liver	Produces bile, which is required for the emulsification and absorption of lipids in the small intestine.
Gallbladder	Stores and concentrates bile and releases it into the small intestine.
Mouth	See the functions of the tongue, salivary glands, and teeth, all of which are in the mouth. Additionally, the lips and cheeks keep food between the teeth during mastication, and buccal glands lining the mouth produce saliva.
Pharynx	Receives a bolus from the oral cavity and passes it into the esophagus.
Esophagus	Receives a bolus from the pharynx and moves it into the stomach; this requires relaxation of the upper esophageal sphincter and secretion of mucus.
Stomach	Mixing waves combine saliva, food, and gastric juice, which activates pepsin, initiates protein digestion, kills microbes in food, helps absorb vitamin B ₁₂ , contracts the lower esophageal sphincter, increases stomach motility, relaxes the pyloric sphincter, and moves chyme into the small intestine.
Small Intestine	Segmentation mixes chyme with digestive juices; peristalsis propels chyme toward the ileocecal sphincter; digestive secretions from the small intestine, pancreas, and liver complete the digestion of carbohydrates, proteins, lipids, and nucleic acids; circular folds, villi, and microvilli help absorb about 90% of digested nutrients.
Large Intestine	Haustral churning, peristalsis, and mass peristalsis drive the colonic contents into the rectum; bacteria produce some B vitamins and vitamin K; absorption of some water, ions, and vitamins occurs; defecation.

Which of the following presents the structures through which chyme travels in the correct sequence?

- A. oesophagus, ileum, duodenum, ileo-cecal valve, transverse colon, rectum
- B. stomach, duodenum, ileum, transverse colon, ileo-cecal valve, rectum
- C. stomach, duodenum, transverse colon, ileum, rectum, ileo-cecal valve,
- D. ileo-cecal valve, stomach, duodenum, transverse colon, ileum, rectum

B

